

GALLIVAC IB88

**Live vaccine against Infectious Bronchitis
caused by a coronavirus variant, strain
CR88121
FREEZE-DRIED PELLETT**

COMPOSITION

Each dose of vaccine contains:

- attenuated virus, strain CR88121, at least 4 log₁₀ EID₅₀
- excipient, q.s. 1 dose

INDICATION

Active immunization of chickens against Infectious Bronchitis caused by the coronavirus variant of group CR88.

ADMINISTRATION AND DOSAGE

Reconstitution of the vaccine:

Using a disposable syringe, inject 3 to 5mL of demineralised or distilled water into the bottle of vaccine. Withdraw the reconstituted vaccine after complete dissolution into the syringe and inject its contents into the container containing the volume of demineralised or distilled water necessary for the administration of the product. In order to recover all the product, rinse the bottle twice.

Route of administration and posology (for 1000 doses)

Nebulization (respiratory route):

Dilute the reconstituted vaccine solution in clean, antiseptic- or disinfectant-free drinking water and equipment. Use the coarse spray application method (*constant pressure sprayer, or type ULVAVAC™*) capable of producing microdroplets (*average droplet diameter of 100 to 150µm*). Each bird must receive the full recommended dose. The amount of water per 1,000 birds will depend on the equipment used. Sprayer producer instructions should be followed. Wherever a ventilation system is operating in the farm, it must be turned off just before initiating the spray vaccination and kept in off position all along the administration and few minutes after its completion.

Ocular route:

Dilute the reconstituted vaccine solution in 30mL of mineral water or ocular diluent. Place a drop (*30µL*) of the vaccine solution on the eye of each bird. Wait until the drop spreads properly then free the bird.

VACCINATION SCHEDULE

- Broilers: one dose to be administered between 10 and 15 days of age
- Pullets:
 - a) one dose to be administered between 10 and 15 days of age to prevent respiratory syndrome
 - b) one dose to be administered between 8 and 12 weeks of age boosted by the inactivated vaccine injected 3-5 weeks prior to the point of lay

CONTRAINDICATION

None

PRECAUTIONS

- Vaccinate only healthy animals
- Apply usual aseptic procedures
- Use clean, antiseptic- and/or disinfectant-free equipment for the preparation of the vaccine solution
- Destroy unused vaccinal contents and disinfect empty bottles before discarding them
- Do not use "atomizer"-type sprayer for the administration of the vaccine
- This vaccination programme do not replace the vaccination programme with the strain Infectious Bronchitis Mass H120

WITHDRAWAL PERIOD

None

STORAGE

Store between +2°C and +8°C, protected from light

FOR VETERINARY USE ONLY